

AN EFFECTIVE VERSION OF KATOK'S HORSESHOE THEOREM FOR CONSERVATIVE C^2 SURFACE DIFFEOMORPHISMS

BASSAM FAYAD AND ZHIYUAN ZHANG

ABSTRACT. For area preserving C^2 surface diffeomorphisms, we give an explicit finite information condition, on the exponential growth of the number of Bowen's (n, δ) -balls needed to cover a positive proportion of the space, that is sufficient to guarantee positive topological entropy. This can be seen as an effective version of Katok's horseshoe theorem in the conservative setting. We also show that the analogous result is false in dimension larger than 3.

1. INTRODUCTION

Let X be a compact smooth surface with a Riemannian metric. Denote by $\text{Diff}_{\text{vol}}^r(X)$ the group of C^r diffeomorphisms which preserve the volume form m induced by the Riemannian metric. Without loss of generality, we assume that $m(X) = 1$.

A well-known result of Katok, based on Pesin theory, says that if $f \in \text{Diff}^{1+\epsilon}(X)$ has non-zero Lyapunov exponent for some f -invariant non atomic measure, then the topological entropy of f is positive and that f actually has invariant horseshoes that carry most of the topological entropy (see for example [5], or [6]). In particular, this is the case for any $f \in \text{Diff}_{\text{vol}}^{1+\epsilon}(X)$ having positive Lyapunov exponents on a positive measure set, or in other words, when f has positive metric entropy by Pesin's formula.

Besides the positivity of Lyapunov exponents, another manifestation of positive metric entropy is the exponential rate of growth of the Bowen (n, δ) -balls (see Definition 1) that are needed to cover a definite proportion of X (see for example [6]).

DEFINITION 1. Given a continuous map $f : X \rightarrow X$. For any $\delta > 0$, integer $n \geq 1$, any $x \in X$, we define Bowen's (n, δ) -ball centered at x by

$$B_f(x, n, \delta) = \{y \mid d(f^i(x), f^i(y)) < \delta, \forall 0 \leq i \leq n-1\}$$

Given an f -invariant measure μ . For any $\epsilon \in (0, 1)$, let $N_f(n, \delta, \epsilon) = \inf_{\mathcal{U}} |\mathcal{U}|$, where $\mathcal{U}$ is taken over all the subsets of $\{B_f(x, n, \delta)\}_{x \in X}$ such that the union of (n, δ) -balls in $\mathcal{U}$ has μ -measure not less than $1 - \epsilon$. For a finite set I , we use $|I|$ to denote the cardinality of I .

Date: March 21, 2017.

By the sub-additive growth of the number of Bowen balls and Katok's horseshoe theorem, the following statement is true by compactity:

Fact: If the C^2 norm of f is bounded by $D > 0$, and if $h, \delta, \varepsilon > 0$ are fixed, then there exists $n_0 = n_0(D, h, \delta, \varepsilon) > 0$ such that if $N_f(n, \delta, \varepsilon) \geq e^{nh}$ for some integer $n > n_0$, then f has positive topological entropy.

Sketch of proof. Assume by contradiction that there exists $h, \delta, \varepsilon > 0$ and a sequence f_n with a uniform bound on its C^2 norm for which $N_{f_n}(n, \delta, \varepsilon) \geq e^{nh}$ and $h_{\text{top}}(f_n) = 0$. By compactity we can, up to passing to a subsequence, assume that f_n has a limit f that is $C^{1+\text{Lip}}$. Since for any g , the minimal number $N_g(n, \delta)$ needed to cover all of X is essentially sub-additive in n , we have that for a fixed $k \in \mathbb{N}$, and for any n sufficiently large $N_{f_n}(k, \delta) \geq e^{kh/2}$. Therefore $N_f(k, \delta) \geq e^{kh/2}$ for any $k \in \mathbb{N}$ and hence f has positive topological entropy. By Katok's horseshoe theorem, this contradicts the assumption $h_{\text{top}}(f_n) = 0$ for all n . $\square$

In this paper, we will give a direct proof of the above fact that also provides an explicit upper bound for $n_0(D, h, \delta, \varepsilon)$. Our bound will essentially be a tower-exponential of height $K \sim \log(\frac{\log A}{h})$ where $A = \|f\|_{C^1}$. The norm of the second derivative of f enters into the argument of the tower-exponential bound. We will not use in our proof *any* ergodic theory.

Our main tool is a finite information closing lemma for a map $g \in \text{Diff}_{\text{vol}}^2(X)$ that generalizes the one obtained in [2, Theorem 4]. Theorem 4 in [2] asserts that if x is such that $\|Dg^q(x)\|$ is comparable to $\|Dg\|^{\theta q}$ where θ is close to 1 and q is sufficiently large compared to powers of the C^2 norm of g , then there exists a hyperbolic periodic point that shadows a piece of a length q orbit of x . A similar effective closing lemma was previously obtained by Climenhaga and Pesin in [4] for $C^{1+\varepsilon}$ -diffeomorphisms in any dimension, assuming however the existence of a splitting of the tangent spaces along a long orbit with some additional estimates of effective hyperbolicity. For an interesting application of the latter effective approach, we refer the reader to [3].

In this note we will need a generalized version of the effective closing lemma in [2] that gives a shadowing of x by a hyperbolic periodic orbit, even when $\|Dg^q(x)\|$ is much smaller than $\|Dg\|^{\theta q}$, provided that $\|Dg^q(x)\| \geq \|Dg(g^i(x))\|^{\theta q}$, for most of the $i \in [0, q]$. An inductive use of this closing lemma allows one to obtain, under the growth condition of the (n, δ) -balls, sufficiently many hyperbolic periodic points with a good control on their local stable and unstable manifolds to insure the existence of a horseshoe. Note that, in order to exploit the growth condition of the Bowen balls, we need sufficiently precise informations from the shadowing property, which are not covered by the direct bootstrapping of Theorem 4 in [2].

With the same approach, we are also able to conclude positive topological entropy from derivative growth at an explicit time scale along a single, yet not too concentrated, orbit.

1.1. Statements of the main results. Throughout this note, X is a compact surface with a volume form m . Without loss of generality, we assume that $m(X) = 1$. We will denote by $f : X \rightarrow X$ a C^2 diffeomorphism that preserves m such that for constants $A, D > 0$,

$$(*) \quad \begin{cases} \|Df\| & \leq A, \\ \|D^2f\| & \leq D. \end{cases}$$

Here $\|Df\|, \|D^2f\|$ denote respectively the supremum of the first and second derivatives of f .

All the constants that appear in the text will implicitly depend on the surface X .

To simplify notations, we define the following.

DEFINITION 2. For $R_0, R_1 > 0, K \in \mathbb{Z}_+$, we define function $Tower : \mathbb{R}_+^2 \times \mathbb{Z}_+ \rightarrow \mathbb{R}$ by the following recurrence relation,

$$(1.1) \quad Tower(R_0, R_1, K) = \begin{cases} R_0, & K = 1, \\ R_1^{Tower(R_0, R_1, K-1)}, & K \geq 2. \end{cases}$$

Our main result is the following.

Theorem A. There exists a constant $C_0 = C_0(X) > 0$ such that the following is true. For any $A, D > 1, h \in (0, \log A], \varepsilon \in (0, 1), \delta > 0$, denote by

$$(1.2) \quad P_0 = \max(\varepsilon^{-1} e^{C_0(\log(\frac{\log A}{h}))^2 + C_0}, C_0 h^{-1} \log \delta^{-1}),$$

$$(1.3) \quad P_1 = e^{C_0 h^{-1} \log D \log A}.$$

If $f : X \rightarrow X$ is a C^2 diffeomorphism preserving m that satisfies $(*)$, and $N_f(n, \delta, \varepsilon) > e^{nh}$ for some $n \geq Tower(P_0, P_1, K_0)$, where $K_0 = \lceil C_0 \log(\frac{\log A}{h}) + C_0 \rceil$, then f has positive topological entropy.

Theorem A gives positive topological entropy from complexity growth at an explicit large time scale. Some adaptation of the proof also allows us to conclude positive topological entropy from derivative growth at an explicit time scale along a single, yet not too concentrated, orbit. To precisely formulate such a result, we introduce the following notation.

DEFINITION 3. Given a continuous map $f : X \rightarrow X$, for any subset $I \subset \mathbb{Z}$, any $x \in X$, we set $Orb(f, x, I) = \{f^i(x) | i \in I\}$.

For constants $c, \delta > 0, \varepsilon \in (0, 1)$, we say that x is $(n, c, \delta, \varepsilon)$ -sparse if for any subset $I \subset \{0, \dots, n-1\}$ satisfying $|I| > cn$ we have $m(B(Orb(f, x, I), \delta)) > \varepsilon$.

Theorem B. There exists a constant $C_0 = C_0(X) > 0$ such that the following is true. For any $A, D > 1$, $h \in (0, \log A]$, $\varepsilon \in (0, 1)$, let

$$P_0 = \varepsilon^{-1} e^{C_0(\log(\frac{\log A}{h}))^2 + C_0}, \quad P_1 = e^{C_0 h^{-1} \log D \log A}.$$

If $f : X \rightarrow X$ is a C^2 diffeomorphism preserving m that satisfies $(*)$, and there exists $x \in X$ such that for some $n \geq \text{Tower}(P_0, P_1, K_0)$, where $K_0 = \lceil C_0 \log(\frac{\log A}{h}) + C_0 \rceil$, we have

- $\|Df^n(x)\| > e^{nh}$,
- x is $(n, \text{Tower}(P_0, P_1, K_0 - 1)^{-1}, D^{-\text{Tower}(P_0, P_1, K_0 - 1)}, \varepsilon)$ -sparse,

then f has positive topological entropy.

Observe that a non-concentration condition, such as the second condition of Theorem B, is necessary to conclude positive entropy, for otherwise x could just belong to a hyperbolic periodic orbit with a small period.

We remark that Theorem A does not hold in general in dimension at least 4 as the following example shows.

Example 1. Denote by $\{g_t\}_{t \in \mathbb{R}}$ a geodesic flow on $Y := S_1 M$, the unit tangent space of a hyperbolic surface M , preserving the Liouville measure μ . We set $h_0 := h_\mu(g_1) > 0$. Let $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ be the circle and let $\varphi \in C^\infty(\mathbb{T})$ be a function such that $\int_{\mathbb{T}} \varphi d\theta = 0$ and $\varphi|_{[0, \frac{1}{2}]} \equiv 1$. For any $\alpha \in \mathbb{R}$, denote by $R_\alpha : \mathbb{T} \rightarrow \mathbb{T}$ the rotation $\theta \mapsto \theta + \alpha[1]$, and consider the C^2 map $f_\alpha : \mathbb{T} \times Y \rightarrow \mathbb{T} \times Y$ defined as follows,

$$f_\alpha(\theta, x) = (\theta + \alpha, g_{\varphi(\theta)}(x)), \quad \forall (\theta, x) \in \mathbb{T} \times Y.$$

Observe that for any $\alpha \in \mathbb{R}$, f_α preserves the smooth measure $\nu := \text{Leb}_{\mathbb{T}} \times \mu$. It is clear that $\sup_{\alpha \in \mathbb{T}} \|f_\alpha\|_{C^2} < \infty$. Moreover, we have the following that shows that Theorem A does not hold in general in dimension at least 4.

PROPOSITION 1. *We have that*

- (1) *For any $\alpha \in \mathbb{R} - \mathbb{Q}$, the topological entropy $h_{\text{top}}(f_\alpha) = 0$.*
- (2) *There exists $\delta > 0$ such that for any $\varepsilon \in (0, 1)$, any integer $n_0 > 0$, there exists $n > n_0$, $\bar{\alpha} \in \mathbb{T}$, such that for any $\alpha \in [0, \bar{\alpha}]$ it holds that $N_{f_\alpha}(n, \delta, \varepsilon) > e^{\frac{nh_0}{2}}$.*

Proof. Abramov Rohlin formula for the entropy of a skew product yields (1) [1]. To see (1) directly, let $(q_n)_{n \in \mathbb{N}}$ be the sequence of denominators of the best rational approximations of α . Then by Denjoy-Koksma theorem, the partial sums $S_{q_n} \varphi$ defined as $S_{q_n} \varphi(\theta) := \sum_{i=0}^{q_n-1} \varphi(\theta + i\alpha)$, $\forall \theta \in \mathbb{T}$, converge uniformly in the C^∞ topology to 0, as n tends to infinity. By direct computations, we see that

$$f_\alpha^{q_n}(\theta, x) = (\theta + q_n \alpha, g_{S_{q_n} \varphi(\theta)}(x)), \quad \forall (\theta, x) \in \mathbb{T} \times Y.$$

This implies that $f_\alpha^{q_n}$ converge to Id in the C^∞ topology, as n tends to infinity. By Ruelle's entropy inequality, such convergence can happen only if $h_{top}(f_\alpha) = 0$.

To see (2), we notice that by $h_\mu(g_1) = h_0 > 0$, there exists $\delta > 0$, such that for any $\varepsilon \in (0, 1)$, any $n_0 > 0$, there exists $n > n_0$ such that $N_{g_1}(n, \delta, \varepsilon) > e^{\frac{nh_0}{2}}$. Then by choosing α to be sufficiently close to 0, so that $i\alpha \in [0, \frac{1}{2}]$ for all $0 \leq i \leq n$, we have $f_\alpha^i(\theta, x) = (\theta + i\alpha, g_i(x))$ for any $(\theta, x) \in \mathbb{T} \times Y$, any $0 \leq i \leq n$. Then it is direct to see that $N_{f_\alpha}(n, \delta, \varepsilon) \geq N_{g_1}(n, \delta, \varepsilon) > e^{\frac{nh_0}{2}}$. This concludes the proof. $\square$

NOTATION 1. For any $n \geq 1$, any $x \in X$, we will denote by $\mu_{x,n} = \frac{1}{n} \sum_{m=0}^{n-1} \delta_{f^m(x)}$. For any $x \in X$, any linear subspace $E \subset T_x X$, any $r > 0$, we denote by $B_E(r) = \{v \in E \mid \|v\| < r\}$. For any subset $A \subset X$, any $r > 0$, we denote by $B(A, r) = \{x \mid d(x, A) < r\}$. For any measurable subset $K \subset X$, we use $|K|$ or $m(K)$ to denote the measure of K .

We will use $c, c_1, \dots$ to denote generic positive constants which are allowed to vary from line to line, and may or may not depend on X , but independent of everything else. Under our notations, expressions like $cA \leq B \leq cA$ are legitimate. For two variables $A, B > 0$, we denote $A \gg B$ (resp. $A \ll B$) if we have $A \geq cB$ (resp. $cA \leq B$) for some constant c as above.

2. FROM HYPERBOLIC POINTS TO POSITIVE ENTROPY

DEFINITION 4. Let $g : X \rightarrow X$ be a C^1 diffeomorphism. For $\alpha \in (0, \pi), r \in (0, 1)$, a hyperbolic periodic point of g , denoted by $y \in X$, is said to be (α, r) -hyperbolic if the following is true. Let $E^s(y), E^u(y)$ be respectively the stable and unstable direction at y . Then

- (1) The angle between $E^s(y)$ and $E^u(y)$ is at least α ,
- (2) The local stable (resp. local unstable) manifold of g at x contains $\exp_y(\text{graph}(\gamma_s))$ (resp. $\exp_y(\text{graph}(\gamma_u))$), where $\gamma_s : B_{E^s(y)}(r) \rightarrow E^u(y)$ (resp. $\gamma_u : B_{E^u(y)}(r) \rightarrow E^s(y)$) is a Lipschitz function such that $\gamma_s(0) = 0$ and $\text{Lip}(\gamma_s) < \frac{1}{100}$ (resp. $\gamma_u(0) = 0$ and $\text{Lip}(\gamma_u) < \frac{1}{100}$).

Moreover, we denote $\exp_y(\text{graph}(\gamma_s))$ (resp. $\exp_y(\text{graph}(\gamma_u))$) by $\mathcal{W}_r^s(y)$ (resp. $\mathcal{W}_r^u(y)$).

For any $\alpha \in (0, \pi), r > 0$, the set of all (α, r) -hyperbolic points of g is denoted by $\mathcal{H}(g, \alpha, r)$. To simplify notations, for any $\lambda \in (0, 1)$, a (λ^2, λ^3) -hyperbolic point of g is said to be λ -hyperbolic. The set of all λ -hyperbolic points of g is denoted by $\mathcal{H}(g, \lambda)$.

DEFINITION 5 (Heteroclinic intersection). For any C^1 diffeomorphism $g : X \rightarrow X$, for any two distinct hyperbolic periodic points of g denoted by p, q , we say that p, q has a heteroclinic intersection, if the stable submanifold of p intersects transversely the unstable manifold of q , and the unstable submanifold of p intersects transversely the stable manifold of q .

The following simple lemma shows that for any given α, r , there cannot be too many (α, r) -points unless there is a heteroclinic intersection.

PROPOSITION 2. *There exist $C_1, C_2 > 1$ depending only on X such that, for any $\alpha \in (0, \pi)$, any $0 < r < C_1^{-1}$, if a C^1 diffeomorphism $g : X \rightarrow X$ satisfy $|\mathcal{H}(g, \alpha, r)| > C_2 r^{-2} \alpha^{-4}$, then there exists a heteroclinic intersection for g . In particular, g has positive topological entropy. In particular, if $\lambda \ll 1$ and $\mathcal{H}(g, \lambda) \gg \lambda^{-14}$, then there exists a heteroclinic intersection for g .*

Proof. In order to be able to measure the angles between vectors in nearby tangent spaces, we cover the surface X by finitely many C^∞ local charts $\{\psi : [-1, 2]^2 \rightarrow X\}_{\psi \in \mathcal{B}}$ indexed by $\mathcal{B}$. For any three distinct points $x, y, z \in \mathbb{R}^2$, let $\angle(x, y, z)$ denote $\angle(x - y, z - y)$. For any $\beta > 0$, any $v \in \mathbb{R}^2 \setminus \{0\}$, let $C(v, \beta) := \{u | \angle(u, v) < \beta\} \cup \{0\}$.

We will choose $\{\psi : [-1, 2]^2 \rightarrow X\}_{\psi \in \mathcal{B}}$ and a constant $c_0 > 0$, depending only on X , such that for any $x \in X$, any $\psi \in \mathcal{B}$ such that $x \in \psi([-1, 1]^2)$, for any $v_1, v_2 \in T_x X \setminus \{0\}$, set $\hat{x} := \psi^{-1}(x)$, $\hat{v}_1 := D\psi^{-1}(x, v_1)$, $\hat{v}_2 := D\psi^{-1}(x, v_2)$, then :

- (1) $2^{-1}\angle(v_1, v_2) \leq \angle(\hat{v}_1, \hat{v}_2) \leq 2\angle(v_1, v_2)$,
 - (2) If $\|v_1\|, \|v_2\| < 2c_0^{-1}$, then $\psi^{-1}\exp_x(v_i)$ is defined and
- $$2^{-1}\angle(v_1, v_2) \leq \angle(\psi^{-1}\exp_x(v_1), \hat{x}, \psi^{-1}\exp_x(v_2)) \leq 2\angle(v_1, v_2).$$

We fix an arbitrary smooth measure $\hat{m}$ on compact manifold

$$\hat{X} = \{(x, v_1, v_2) | x \in X, v_1, v_2 \in T_x X, \|v_1\| = \|v_2\| = c_0^{-1}\}.$$

Let $c_1 > 0$ be a large constant to determined later, and for any $(x, v_1, v_2) \in \hat{X}$, any $\psi \in \mathcal{B}$ so that $x \in \psi((0, 1)^2)$ and set

$$Q_\psi(x, v_1, v_2) = \{(y, u_1, u_2) \in \hat{X} | |\hat{x} - \hat{y}| < \frac{r\alpha}{c_1}, \angle(\hat{v}_1, \hat{u}_1), \angle(\hat{v}_2, \hat{u}_2) < \frac{\alpha}{40}\}.$$

Then there exists $c_2 > 0$ depending only on X, c_1 , such that for all $(x, v_1, v_2) \in \hat{X}$, any $\psi \in \mathcal{B}$ so that $x \in \psi((0, 1)^2)$, we have

$$\hat{m}(\psi(Q_\psi(x, v_1, v_2))) > c_2^{-1} r^2 \alpha^4.$$

By pigeonhole principle, there exists a constant $c_3 > 0$ depending only on X, c_2 , such that whenever $|\mathcal{H}(g, \alpha, r)| > c_3 r^{-2} \alpha^{-4}$, there exists a chart $\psi \in \mathcal{B}$, $(y_i, v_i^s, v_i^u) \in \hat{X}$, $i = 1, 2$ so that

- (1) $y_1, y_2 \in \mathcal{H}(g, \alpha, r) \cap \psi((0, 1)^2)$ are two distinct points;
- (2) for $i = 1, 2$, $\angle(v_i^s, v_i^u) \leq \frac{\pi}{2}$, and v_i^s (resp. v_i^u) lies in the stable (resp. unstable) direction of y_i ;
- (3) $Q_\psi(y_1, v_1^s, v_1^u) \cap Q_\psi(y_2, v_2^s, v_2^u) \neq \emptyset$.

This implies that $|\hat{y}_1 - \hat{y}_2| < \frac{2r\alpha}{c_1}$, $\angle(\hat{v}_1^s, \hat{v}_2^s) < \frac{\alpha}{20}$ and $\angle(\hat{v}_1^u, \hat{v}_2^u) < \frac{\alpha}{20}$.

For $i = 1, 2$, let us denote $\alpha_i = \angle(v_i^u, v_i^s)$. By the definition of $\mathcal{H}(g, \alpha, r)$ we have $\alpha_1, \alpha_2 \geq \alpha$. Then $\angle(\hat{v}_i^u, \hat{v}_i^s) \geq 2^{-1}\alpha_i$ for $i = 1, 2$. Moreover for $r \ll 1$, we have $\psi^{-1}(\mathcal{W}_r^u(y_i)) \subset \hat{y}_i + C(\hat{v}_i^u, \frac{1}{20}\alpha_i)$ since there exists $\gamma_u :$

$B_{E^u(y_i)}(r) \rightarrow E^s(y_i)$ with $Lip(\gamma_u) < \frac{1}{100}$, such that $\mathcal{W}_r^u(y_i) = \exp_{y_i} \text{graph}(\gamma_u)$ and $\text{graph}(\gamma_u) \subset C(v_i^u, \frac{1}{40}\alpha_i)$. Similarly, we have $\psi^{-1}(\mathcal{W}_r^s(y_i)) \subset \hat{y}_i + C(\hat{v}_i^s, \frac{1}{20}\alpha_i)$.

By straightforward calculations, when c_1 is chosen to be sufficiently large, y_1, y_2 above have a heteroclinic intersection. Thus for any $r \ll 1$, any C^1 diffeomorphism $g : X \rightarrow X$ so that $|\mathcal{H}(g, \alpha, r)| \gg r^{-2}\alpha^{-4}$, there exists a heteroclinic intersection for g . It is a standard fact that for C^1 surface diffeomorphism, the existence of a heteroclinic intersection implies positive topological entropy. This concludes the proof. $\square$

3. A CLOSING LEMMA

DEFINITION 6. For any $\eta > 0$, any integer $l > 0$, any C^0 map $g : X \rightarrow X$, any subset $Y \subset X$, a point $x \in X$ is said to be (η, l, g) -recurrent for Y if we have

$$\frac{1}{l} |\{0 \leq j \leq l-1 \mid g^j(x) \in Y\}| > \eta.$$

For any subset $Y \subset X$, we denote by

$$\mathcal{R}(Y, \eta, l, g) := \{(\eta, l, g) - \text{recurrent points for } Y\}.$$

For any $\lambda, \xi > 0$, we set

$$(3.1) \quad \mathcal{G}(\lambda, \xi, g) := \bigcup_{y \in \mathcal{H}(g, \lambda)} B(\mathcal{W}_{\lambda^3}^u(y), \xi).$$

By our definition, we clearly have $\mathcal{G}(\lambda, \xi, g) = \mathcal{G}(\lambda, \xi, g^k)$ for any $k \geq 1$, since $\mathcal{H}(g, \lambda) = \mathcal{H}(g^k, \lambda)$ for any $k \geq 1$.

Theorem 4 in [2] can be strengthened to prove the following proposition.

PROPOSITION 3. *There exist $C = C(X) > 1$, and an absolute constant $\theta_0 \in (\frac{1}{2}, 1)$ such that the following is true. For each $\Delta \geq 1$, we set*

$$(3.2) \quad \eta = \eta(\Delta) := C^{-1}\Delta^{-2} \in (0, 1)$$

Let $g : X \rightarrow X$ be a C^2 diffeomorphism preserving m . If for $A_1 \geq C$, $D_1 \geq A_1$, an integer $q \geq D_1^{C\Delta}$ and $x \in X$, we have the following :

- (1) $\|Dg\| \leq A_1^\Delta$,
- (2) $\|D^2g\| \leq D_1$,
- (3) $x \notin \mathcal{R}(\{y \mid \|Dg(y)\| > A_1^{\theta_0^{-1}}\}, \eta, q, g)$,
- (4) $\|Dg^q(x)\| > A_1^q$,

then

$$x \in \mathcal{F}(A_1, D_1, \Delta, q, g) := \bigcup_{1 \leq j \leq q} g^{-j}(\mathcal{G}(D_1^{-C\Delta}, A_1^{-\frac{q}{2D_1^{C\Delta}}}, g))$$

The proof of Proposition 3 follows closely that of Theorem 4 in [2]. In our case we need to get more precise informations on the regularity of local invariant manifolds, as well as the location of the hyperbolic point. We defer its proof to Appendix A relying on many estimates from [2].

4. ESTIMATES ALONG A TOWER EXPONENTIAL SEQUENCE

Without loss of generality, we will always assume that D, A in Theorem A, B satisfy

$$(4.1) \quad D > A \gg 1.$$

Then we can assume that for any C^2 map $g : X \rightarrow X$ such that $\|Dg\|, \|D^2g\| \leq D$, we have

$$\|D^2g^k\| < D^k, \quad \forall k \geq 1.$$

Let C, θ_0 be defined in Proposition 3. For D, A, h given in Theorem A or B, set C' to be a large positive constant depending only on X to be determined later. We set

$$(4.2) \quad \Delta = \frac{16 \log A}{h}, \quad K = \lceil \frac{\log(\frac{\Delta}{4})}{-\log \theta_0} \rceil \geq 2, \quad \eta = \eta(\Delta) \text{ (see (3.2)).}$$

Define

$$(4.3) \quad H = H(X, A, h) := C' \Delta.$$

Given an integer $n \geq 1, \varepsilon \in (0, 1)$, we inductively define the following.

$$(4.4) \quad q_0 = \lceil \varepsilon^{-1} e^{C'(\log \Delta)^2} \rceil,$$

$$(4.5) \quad l_k = \begin{cases} \lceil D^{q_k H} \rceil, & 0 \leq k \leq K-1 \\ \lceil \frac{n}{q_K} \rceil, & k = K \end{cases}, \quad q_{k+1} = q_k l_k.$$

For $0 \leq k \leq K$, we set

$$(4.6) \quad \lambda_k = D^{-C \Delta q_k}, \quad \xi_k = A^{-\frac{q_{k+1} \theta_0^{k+1}}{2D^{C \Delta q_k}}},$$

and set

$$Q_0 = \varepsilon^{-1} e^{C'(\log \Delta)^2}, \quad Q_1 = e^{20C'h^{-1} \log D \log A}.$$

We have the following simple lemma.

- LEMMA 1. (1) $e^{\frac{h}{16}} < A^{\theta_0^{K+1}} \leq e^{\frac{h}{4}}$,
 (2) For any $C' \gg 1$, for all $0 \leq k \leq K-1$, we have $D^{q_k H} \leq l_k \leq \text{Tower}(Q_0, Q_1, k+2)$. If $n > \text{Tower}(Q_0, Q_1, K+3)$, then $l_K \geq D^{q_K H}$,
 (3) For any $C' \gg 1$, set $\delta_0 = D^{-\text{Tower}(Q_0, Q_1, K+1)}$, we have

$$\begin{aligned} \xi_i &\leq \xi_0, \quad \delta_0 < \min(\lambda_K^3, \xi_0), \\ C' \lambda_i^{-11} \max(\delta_0, \xi_i) &< \varepsilon, \quad \forall 0 \leq i \leq K. \end{aligned}$$

We define for $0 \leq k \leq K$,

$$(4.7) \quad \mathcal{G}_k := \mathcal{G}(\lambda_k, \xi_k, f),$$

$$(4.8) \quad \mathcal{F}_k := \mathcal{F}(A^{q_k \theta_0^{k+1}}, D^{q_k}, \Delta, l_k, f^{q_k}).$$

The following is a corollary of Proposition 3.

COROLLARY A. *If $n > \text{Tower}(Q_0, Q_1, K+3)$, then for any $0 \leq k \leq K$ we have*

$$x \notin \mathcal{R}(\{y \mid \|Df^{q_k}(y)\| > A^{q_k \theta_0^k}\}, \eta(\Delta), l_k, f^{q_k}) \cup \mathcal{F}_k \implies \|Df^{q_{k+1}}(x)\| \leq A^{q_{k+1} \theta_0^{k+1}}.$$

Proof. By Lemma 1(2), if $n > \text{Tower}(Q_0, Q_1, K+3)$ then for any $0 \leq k \leq K$, we have $l_K \geq D^{q_k H}$. By our choice of A, D , we have

$$\|Df^{q_k}\| \leq A^{q_k}, \quad \|D^2 f^{q_k}\| < D^{q_k}, \quad \forall 0 \leq k \leq K.$$

We take any $0 \leq k \leq K$, and an arbitrary point $x \in X$ such that $\|Df^{q_{k+1}}(x)\| > A^{q_{k+1} \theta_0^{k+1}}$. It suffices to show that $x \in \mathcal{R}(\{y \mid \|Df^{q_k}(y)\| > A^{q_k \theta_0^k}\}, \eta(\Delta), l_k, f^{q_k}) \cup \mathcal{F}_k$.

By Lemma 1(1) we have $\|Df^{q_k}(x)\| \leq A^{q_k} \leq (A^{q_k \theta_0^{k+1}})^{\frac{16 \log A}{h}}$. If $x \in \mathcal{R}(\{y \mid \|Df^{q_k}(y)\| > A^{q_k \theta_0^k}\}, \eta(\Delta), l_k, f^{q_k})$, we are done. Otherwise, we can verify conditions (1)-(4) in Proposition 3 for $(f^{q_k}, A^{q_k \theta_0^{k+1}}, D^{q_k}, \frac{16 \log A}{h}, l_k)$ in place of (g, A_1, D_1, Δ, q) . We can apply Proposition 3 for map $g = f^{q_k}$ to show that $x \in \mathcal{F}_k$. This completes the proof. $\square$

The following is a straightforward consequence of Proposition 2.

COROLLARY B. *For all $C' \gg 1$ the following is true. If we have at least one of the following :*

- (1) *there exists $0 \leq i \leq K$ such that $|\mathcal{G}_i| \geq \frac{\eta^{K-i} \varepsilon}{(K+1)l_i}$,*
- (2) *there exists $0 \leq i \leq K-1$ such that $m(B(\mathcal{G}_i, D^{-\text{Tower}(Q_0, Q_1, K+3)})) > \varepsilon$,*
then f has a heteroclinic intersection, in which case f has positive topological entropy.

We include the proof of Corollary B in Appendix B.

REMARK 1. *Given A, D, h as in Theorem A or B, we will choose C' to be sufficiently large so that the conclusions of both Lemma 1 and Corollary B hold.*

5. AN ITERATIVE DECOMPOSITION

Now we say a few words about the general strategy behind the proof of Theorem A and Theorem B. We will inductively define a sequence of decompositions of the surface X , denoted by $X = M_i \sqcup E_i$. To start the induction, we define $M_0 = X$ and $E_0 = \emptyset$. Assume that for $k \geq 0$, we have defined M_k, E_k satisfying the following condition:

$$\text{For each } x \in M_k, \text{ we have } \|Df^{q_k}(x)\| \leq A^{q_k \theta_0^k}.$$

Then E_{k+1} is defined as the set of the points that up till some finite time scale, either run into E_k with frequency $\geq \eta$, or is shadowed by hyperbolic orbits (of course the first case does not happen if E_k is empty). We

will use Proposition 3 to show that the complement of E_{k+1} , defined as M_{k+1} , again satisfies the induction hypothesis. We then argue that after roughly $K = O(\log(\frac{\log A}{h}))$ steps, E_{K+1} has to be large. This will show that at some previous time scale, there are enough different hyperbolic hyperbolic points to create a homoclinic intersection.

The formal construction is the following. For all $0 \leq k \leq K+1$, we define M_k, E_k through the following inductive formula. Let

$$(5.1) \quad E_0 = \emptyset, \quad M_0 = X$$

and for all $0 \leq k \leq K$, we define

$$(5.2) \quad E_{k+1} = \mathcal{R}(E_k, \eta, l_k, f^{q_k}) \bigcup \mathcal{F}_k,$$

$$(5.3) \quad M_{k+1} = X \setminus E_{k+1}$$

LEMMA 2. *If $n > \text{Tower}(Q_0, Q_1, K+3)$, then for any $0 \leq k \leq K+1$ we have*

$$x \in M_k \implies \|Df^{q_k}(x)\| \leq A^{q_k \theta_0^k}.$$

Proof. This is clear when $k = 0$ by $\|Df\| \leq A$ and sub-multiplicativity. Assume that the lemma is valid for some integer $k \in \{0 \dots, K\}$, then $\{x \mid \|Df^{q_k}(x)\| > A^{q_k \theta_0^k}\} \subset E_k$ (we consider the inclusion valid if both sides are empty). By Corollary A and (5.2), we see that any $x \in X$ such that $\|Df^{q_{k+1}}(x)\| > A^{q_{k+1} \theta_0^{k+1}}$ is contained in E_{k+1} . This completes the induction, thus finishes the proof. $\square$

We will give the proof of Theorem A and B in the next two subsections. In the following, we let C, θ_0 be defined in Proposition 3, let $A, D, h > 0$ be given by Theorem A or B, and let C' be sufficiently large depending only on X , satisfying Remark 1.

5.1. Proof of Theorem A.

PROPOSITION 4. *Let C_0 in Theorem A be sufficiently large. Then under the conditions of Theorem A, we have*

$$|E_{K+1}| \geq \varepsilon.$$

Proof. We first show the following lemma.

LEMMA 3. *Let C_0 in Theorem A be sufficiently large, and let n be given as in Theorem A. Then for each $y \in M_{K+1}$, we have*

$$B(y, e^{-2nh/5} \delta) \subset B_f(y, n, \delta).$$

Proof. It is clear from (4.5) that

$$l_K \in \left(\frac{n}{q_K}, \frac{21}{20} \frac{n}{q_K}\right).$$

Let $y \in M_{K+1}$. For each $0 \leq i \leq l_K - 1$, we denote by

$$a_i = \log \|Df^{q_K}(f^{iq_K}(y))\|, \quad \delta_i = e^{-2nh/5 + iq_K h/24 + \sum_{j=0}^{i-1} a_j} \delta, \quad B_i = B(f^{iq_K}(y), \delta_i).$$

By letting C_0 in Theorem A be sufficiently large, we can ensure that $n > \text{Tower}(P_0, P_1, K_0) > \text{Tower}(Q_0, Q_1, K + 3)$. Then by Lemma 1(1) and Lemma 2, we have for each $z \in M_K$, $\log \|Df^{q_K}(z)\| \leq q_K \theta_0^K \log A \leq \frac{hq_K}{4}$. Then by $y \in M_{K+1}$, (5.2) and Lemma 2, we have $y \notin \mathcal{R}(\{z \mid \log \|Df^{q_K}(z)\| > \frac{hq_K}{4}\}, \eta, l_K, f^{q_K})$, thus

$$|\{0 \leq i \leq l_K - 1 \mid a_i > \frac{hq_K}{4}\}| \leq \eta l_K.$$

Since $0 \leq a_i \leq q_K \log A$ for any $0 \leq i \leq l_K - 1$, we have

$$\sum_{j=0}^{i-1} a_j \leq \sum_{j=0}^{l_K-1} a_j \leq \eta l_K q_K \log A + \frac{l_K q_K h}{4} \leq \frac{7l_K q_K h}{24}, \quad \forall 0 \leq i \leq l_K - 1.$$

The last inequality follows from $\eta \leq \frac{h}{24 \log A}$ which is a consequence of (4.2), (3.2) and $h \in (0, \log A]$. Then for any $0 \leq i \leq l_K - 1$, we have

$$(5.4) \quad \delta_i \leq e^{-2nh/5 + l_K q_K h/3} \delta \leq e^{-\frac{1}{20}nh} \delta.$$

We claim that for any integer $0 \leq i \leq l_K - 1$,

$$(5.5) \quad f^{iq_K}(B_0) \subset B_i.$$

We first show that the above claim concludes the proof of our lemma. Indeed, for any $0 \leq l \leq n$, there exist $0 \leq i \leq l_K - 1, 0 \leq j \leq q_K - 1$ such that $l = iq_K + j$. Then we have

$$f^l(B_0) = f^j(f^{iq_K}(B_0)) \subset f^j(B_i) \subset B(f^l(y), \delta)$$

The last inclusion follows from $A^{q_K} \delta_i \leq A^{q_K} e^{-nh/20} \delta \leq \delta$, by $\|Df^j\| \leq A^{q_K}$, (5.4) and $\frac{n}{q_K} \geq \frac{20 \log D}{h}$.

Now we obviously have (5.5) for $i = 0$. Assume that we have (5.5) for some $0 \leq i \leq l_K - 1$, we will show that we have (5.5) for $i + 1$. It suffices to show that $f^{q_K}(B_i) \subset B_{i+1}$. Using the C^2 bound $\|D^2 f^{q_K}\| \leq D^{q_K}$ and $\frac{n}{q_K} \geq \frac{20 \log D}{h}$, we see that for any $z \in B_i$,

$$\begin{aligned} \|Df^{q_K}(z)\| &\leq e^{a_i} + \delta_i D^{q_K} \\ &\leq e^{a_i} + D^{q_K} e^{-nh/20} \delta \leq e^{a_i + hq_K/24}. \end{aligned}$$

Since $\delta_{i+1} = e^{a_i + hq_K/24} \delta_i$, we obtain $f^{q_K+1}(B_i) \subset B_{i+1}$. This proves (5.5) and concludes the proof of Lemma 3. $\square$

To proceed with the proof of Proposition 4, observe that by Lemma 3, $M_{K+1} = X \setminus E_{K+1}$ can be covered by $ce^{4nh/5} \delta^{-2}$ many Bowen's (n, δ) -balls. By (1.2), $n > P_0$ and by letting C_0 be large, we have $c\delta^{-2} < e^{P_0 h/5} < e^{nh/5}$. This implies that $|E_{K+1}| \geq \varepsilon$. $\square$

Proof of Theorem A. Since f is area preserving, by Markov's inequality we have

$$|\mathcal{R}(E_k, \eta, l_k, f^{q_K})| \leq \eta^{-1} |E_k|.$$

Again by the fact that f is area preserving, we obtain the following inequality by (5.2), (4.8)

$$(5.6) \quad |E_{k+1}| \leq \eta^{-1}|E_k| + |\mathcal{F}_k| \leq \eta^{-1}|E_k| + l_k|\mathcal{G}_k|.$$

By (5.6) and (5.1), we have

$$|E_{K+1}| \leq \sum_{i=0}^K \eta^{i-K} l_i |\mathcal{G}_i|.$$

Thus $|E_{K+1}| \geq \varepsilon$ implies that $|\mathcal{G}_i| \geq \eta^{K-i} \frac{\varepsilon}{(K+1)l_i}$ for some $0 \leq i \leq K$, which by Corollary B (1) implies that f has positive entropy. $\square$

5.2. Proof of Theorem B. The proof of Theorem B is parallel to that of Theorem A. The following proposition is an analogue of Proposition 4.

PROPOSITION 5. *Let C_0 in Theorem B be sufficiently large, and let n be as in Theorem B. Then under the condition of Theorem B, we have*

$$\mu_{x,n}(E_K) \geq \frac{h}{2 \log A}.$$

Proof. By letting C_0 in Theorem A be sufficiently large, we can ensure that $n > \text{Tower}(P_0, P_1, K_0) > \text{Tower}(Q_0, Q_1, K+3)$. Then by Lemma 2, for each $y \in M_K$, we have $\|Df^{q_K}(y)\| \leq A^{q_K \theta_0^K} \leq e^{\frac{q_K h}{2}}$.

We take a subset $\{p_1, \dots, p_l\} \subset \{0, \dots, n - q_K\}$ so that $I_j := \{p_j, \dots, p_j + q_K - 1\}$, $1 \leq j \leq l$ are disjoint subsets of $\{0, \dots, n - 1\}$ and $f^{p_j}(x) \in M_K$ for all j . Moreover, we assume that for any $k \in \{0, \dots, n - 1\} \setminus \bigcup_{j=1}^l I_j$, we have $f^k(x) \in E_K$. The construction of $\{p_i\}_{i=1}^l$ is straightforward.

Then by sub-multiplicativity, we have

$$\begin{aligned} \log \|Df^n(x)\| &\leq \sum_{i=1}^l \log \|Df^{q_K}(f^{p_i}(x))\| + (n - lq_K) \log A \\ &\leq \frac{1}{2} l h q_K + (n - lq_K) \log A. \end{aligned}$$

By the condition in Theorem B, we have $\log \|Df^n(x)\| > nh$. Thus $n(\log A - h) > lq_K(\log A - \frac{h}{2})$. Then we see that $\mu_{x,n}(E_K) \geq \frac{n-lq_K}{n} \geq \frac{h}{2 \log A}$. $\square$

Proof of Theorem B. For any measurable set $B \subset X$, any integers $n, l \geq 1$, any $x \in X$, we have

$$\mu_{x,n}(f^{-l}(B)) \leq \frac{l}{n} + \mu_{x,n}(B).$$

Then for any $k = 0, \dots, K-1$, by Markov's inequality we have

$$\begin{aligned} \mu_{x,n}(\mathcal{R}(E_k, \eta, l_k, f^{q_k})) &\leq (\eta l_k)^{-1} \int \sum_{i=0}^{l_k-1} 1_{f^{-iq_k}(E_k)} d\mu_{x,n} \\ &\leq (\eta l_k)^{-1} \sum_{i=0}^{l_k-1} \mu_{x,n}(f^{-iq_k}(E_k)) \\ &\leq (\eta l_k)^{-1} \sum_{i=0}^{l_k-1} (\mu_{x,n}(E_k) + \frac{iq_k}{n}) \leq \eta^{-1} \mu_{x,n}(E_k) + \frac{q_{k+1}}{2n\eta}. \end{aligned}$$

Similarly, we have

$$\begin{aligned} \mu_{x,n}(\mathcal{F}_k) &\leq \sum_{i=0}^{l_k-1} \mu_{x,n}(g^{-iq_k}(\mathcal{G}_k)) \\ &\leq l_k \mu_{x,n}(\mathcal{G}_k) + \frac{l_k q_{k+1}}{n}. \end{aligned}$$

Then we have an inequality analogous to (5.6), as follows,

$$\begin{aligned} \mu_{x,n}(E_{k+1}) &\leq \mu_{x,n}(\mathcal{R}(E_k, \eta, l_k, f^{q_k})) + \mu_{x,n}(\mathcal{F}_k) \\ &\leq \eta^{-1} \mu_{x,n}(E_k) + \frac{q_{k+1}}{2n\eta} + l_k \mu_{x,n}(\mathcal{G}_k) + \frac{l_k q_{k+1}}{n}. \end{aligned}$$

By the simple observation that $l_k \geq l_0 \geq \eta^{-1}$ for all $0 \leq k \leq K$, we have

$$\mu_{x,n}(E_K) \leq \sum_{i=0}^{K-1} \eta^{-K+i+1} (l_i \mu_{x,n}(\mathcal{G}_i) + \frac{2l_i q_{i+1}}{n}).$$

By (4.5) and Proposition 5, we see that there exists $0 \leq i \leq K-1$ such that

$$\mu_{x,n}(\mathcal{G}_i) \geq l_i^{-1} \left(\frac{\eta^{K-i-1}}{K} \frac{h}{2 \log A} - \frac{2q_K l_{K-1}}{n} \right) \geq l_i^{-2}.$$

The last inequality follows from

$$\begin{aligned} \frac{2q_K l_{K-1}}{n} &< \frac{2l_{K-1}}{l_K} < q_0^{-1} \leq e^{-C'(\log \Delta)^2} < \frac{\eta^K h}{4K \log A}, \\ \frac{\eta^K h}{4K \log A} &> l_0^{-1} \geq l_i^{-1}, \quad \forall 0 \leq i \leq K-1, \end{aligned}$$

by letting C' be larger than some absolute constant. In particular, by Lemma 1(2), (4.2), (4.4), (4.5), and by letting C_0 in Theorem B be sufficiently large, we have

$$\begin{aligned} \mu_{x,n}(\mathcal{G}_i) &> \text{Tower}(Q_0, Q_1, K+1)^{-2} > \text{Tower}(Q_0, Q_1, K+2)^{-1}, \\ K_0 &\geq K+4, \quad P_i > Q_i, i = 0, 1. \end{aligned}$$

By the condition of Theorem B that x is $(n, \text{Tower}(P_0, P_1, K_0-1)^{-1}, D^{-\text{Tower}(P_0, P_1, K_0-1)}, \varepsilon)$ -sparse, we see that

$$m(B(\mathcal{G}_i, D^{-\text{Tower}(Q_0, Q_1, K+3)})) \geq m(B(\mathcal{G}_i, D^{-\text{Tower}(P_0, P_1, K_0-1)})) > \varepsilon.$$

This concludes the proof by Corollary B (2). □

APPENDIX A.

In this section we prove the main technical result Proposition 3. We start with a slight generalization of Pliss lemma [7].

LEMMA 4 (a variant of Pliss). *For any real numbers $N \geq 1$, $1 > \theta_0 > \theta_1 > \theta_2 > 0$, $\eta \in (0, \frac{1}{2} \frac{1-\theta_0}{N-\theta_0} \frac{\theta_1-\theta_2}{N-\theta_2})$, for any integer $n \geq 1$, real number $l > 0$, the following is true. Given a sequence of n real numbers $a_1, \dots, a_n$. Assume that*

- (1) $a_i \leq Nl$ for all $1 \leq i \leq n$,
- (2) $\sum_{i=1}^n a_i > n\theta_1 l$,
- (3) $|\{1 \leq i \leq n \mid a_i > \theta_0 l\}| < \eta n$.

Then there exist at least $\frac{\theta_1-\theta_2}{1-\theta_2} n$ indexes i 's such that $\frac{1}{k} \sum_{j=i}^{i+k-1} a_j > \theta_2 l$ for all $1 \leq k \leq n+1-i$.

Proof. Denote by

$$A := \{i \mid \text{there exists } 1 \leq k \leq n+1-i \text{ such that } \frac{1}{k} \sum_{j=i}^{i+k-1} a_j \leq \theta_2 l\}$$

Without loss of generality, we assume that $A \neq \emptyset$, for otherwise the conclusion of Lemma 4 is already true. Then A is contained in a non-empty set $I \subset \{1, \dots, n\}$ satisfying that $\frac{1}{|I|} \sum_{i \in I} a_i \leq \theta_2 l$. Then by (1),(2), we obtain that

$$lN|I^c| + l\theta_2|I| > l n\theta_1.$$

By $l > 0$, the above inequality implies that $|I^c| \geq \frac{\theta_1-\theta_2}{N-\theta_2} n$. We claim that

$$(A.1) \quad \frac{1}{|I^c|} \sum_{i \in I^c} a_i \leq l.$$

Indeed, if (A.1) was false, by (1) we would have at least $\frac{1-\theta_0}{N-\theta_0} |I^c| \geq \frac{1-\theta_0}{N-\theta_0} \frac{\theta_1-\theta_2}{N-\theta_2} n > \eta n$ indexes $i \in I^c$ such that $a_i > \theta_0 l$, but this would contradict (3).

Now we use (2) again, with the improved estimate (A.1) in place of (1), and obtain

$$l|I^c| + \theta_2 l|I| \geq \sum_{i \in I^c} a_i + \sum_{i \in I} a_i > n\theta_1 l.$$

This implies that $|I^c| \geq \frac{\theta_1-\theta_2}{1-\theta_2} n$. We conclude the proof by the definition of I . □

Let x be given by the condition of Proposition 3. We will define a collection of charts along a sub-orbit of x following the definitions and estimates in [2].

Let v_s be a unit vector in the most contracting direction of $Dg^q(x)$ in $T_x X$, and let v_u be a unit vector orthogonal to v_s . For each $0 \leq i \leq q$, we define

$$\begin{aligned} v_i^s &:= \frac{Dg^i(v_s)}{\|Dg^i(v_s)\|}, & v_i^u &:= \frac{Dg^i(v_u)}{\|Dg^i(v_u)\|}, \\ \lambda_i^s &:= \log \frac{\|Dg(v_i^s)\|}{\|v_i^s\|}, & \lambda_i^u &:= \log \frac{\|Dg(v_i^u)\|}{\|v_i^u\|}, \\ \bar{\lambda}_i^e &:= \min\{\lambda_i^u, -\lambda_i^s\}. \end{aligned}$$

Given $r > 0, \tau > 0, \kappa > 0$, we define a (r, τ, κ) -Box, which we denote by $U(r, \tau, \kappa)$, to be

$$U(r, \tau, \kappa) = \{(v, w) \in \mathbb{R}^2 \mid \|v\| \leq r, \|w\| \leq \tau + \kappa\|v\|\}.$$

For $\kappa > 0$, we denote by

$$\begin{aligned} C(\kappa) &= \{(v, w) \in \mathbb{R}^2 \mid \|w\| < \kappa\|v\|\}, \\ \tilde{C}(\kappa) &= \{(v, w) \in \mathbb{R}^2 \mid \|v\| < \kappa\|w\|\}. \end{aligned}$$

We will refer to these sets as cones.

We now recall some definitions in [2].

- A curve contained in $\mathbb{R}^2 = \mathbb{R}_x \oplus \mathbb{R}_y$ is called a κ -horizontal graph if it is the graph of a Lipschitz function from an closed interval $I \subset \mathbb{R}_x$ to $\mathbb{R}_y$ with Lipschitz constant less than κ . Similarly, we can define the κ -vertical graphs.
- The boundary of an (r, τ, κ) -Box U is the union of two 0-vertical graphs and two κ -horizontal graphs. We call these graphs respectively, the *left (resp. right) vertical boundary of U* and the *upper (resp. lower) horizontal boundary of U* . We call the union of the left and right vertical boundary of U the *vertical boundary of U* . Similarly we call the union of the upper and lower horizontal boundary of U the *horizontal boundary of U* .
- Horizontal and vertical graphs which connect the boundaries of U will be called full horizontal and full vertical graphs as in the following definition. Given $r, \tau, \kappa, \eta > 0$, for each (r, τ, κ) -Box U , an η -full horizontal graph of U is an η -horizontal graph L such that $L \subset U$ and the endpoints of L are contained in the vertical boundary of U . Similarly, we define the η -full vertical graphs of U .
- We define an η -horizontal strip of U to be a subset of U bounded by the vertical boundary of U and two disjoint η -full horizontal graphs of U which are both disjoint from the horizontal boundary of U . Similarly we can define η -vertical strips of U . Like Boxes, we define the horizontal, vertical boundary of a strip.
- Given a Box U , $\mathcal{R}'$ a vertical strip of U , and $\mathcal{R}$ a horizontal strip of U , a homeomorphism that maps $\mathcal{R}'$ to $\mathcal{R}$ is said to be *regular* if it maps the horizontal (resp. vertical) boundary of $\mathcal{R}'$ homeomorphically to the horizontal (resp. vertical) boundary of $\mathcal{R}$.

We recall the definition of hyperbolic map in [2].

DEFINITION 7. Given $r, \tau > 0, 0 < \kappa, \kappa', \kappa'' < 1$. Denote $U = U(r, \tau, \kappa)$, and let $\mathcal{R}_1$ be a κ -vertical strip of U , $\mathcal{R}_2$ be a κ -horizontal strip of U . A C^1 diffeomorphism $G : \mathcal{R}_1 \rightarrow \mathbb{R}^2$ is called a hyperbolic map if it satisfies the following conditions:

(A.2) G is a regular map from $\mathcal{R}_1$ to $\mathcal{R}_2$,

(A.3) $\forall x \in \mathcal{R}_1, DG_x(C(\kappa')) \subset C(\frac{1}{2}\kappa')$,

(A.4) $\forall x \in \mathcal{R}_2, DG_x^{-1}(\tilde{C}(\kappa'')) \subset \tilde{C}(\frac{1}{2}\kappa'')$.

The following is a sketch of a hyperbolic map.




FIGURE 1. $\mathcal{R}_1$ is the topological rectangle $abcd$, $\mathcal{R}_2$ is the topological rectangle $a'b'c'd'$. Under a hyperbolic map G , ab is mapped to $a'b'$. Similarly, bc, cd, da are mapped respectively to $b'c', c'd', d'a'$.

For each $0 \leq n \leq q$, we define $i_n : \mathbb{R}^2 \rightarrow T_{x_n} S$ as

$$i_n(a, b) = av_n^u + bv_n^s.$$

There exists a constant $R = R(X) > 0$ such that exp_{x_n} is a diffeomorphism restricted to $i_n(B(0, D_1^{-\Delta} R))$ and $exp_{x_{n+1}}^{-1}$ is a diffeomorphism restricted to

$g \exp_{x_n} i_n(B(0, D_1^{-\Delta} R))$. Denote by g_n the C^2 diffeomorphism

$$\begin{aligned} g_n : B(0, D_1^{-\Delta} R) &\rightarrow \mathbb{R}^2 \\ g_n(v, w) &= i_{n+1}^{-1} \exp_{x_{n+1}}^{-1} g \exp_{x_n} i_n(v, w). \end{aligned}$$

We set $M := 1000$, and

$$\bar{r} = D_1^{-3\Delta M}, \quad \bar{\kappa} = D_1^{-\Delta M}, \quad \delta = \frac{\log A_1}{100}.$$

The main estimates in [2] are summarised in the following proposition.

PROPOSITION 6. *Under the conditions of Proposition 3 for some absolute constant $\theta_0 \in (0, 1)$ sufficiently close to 1, and $C > 0$ sufficiently large depending only on X , there exist constant $C_1 = C_1(X)$, integers $0 \leq i_1 \leq i_2 \leq q$, and sequences of positive numbers $\{(r_n, \tau_n, \kappa_n, \tilde{\kappa})_n\}_{i_1 \leq n \leq i_2}$ such that :*

(1) (Positive proportion)

$$i_2 - i_1 \geq D_1^{-C_1 \Delta} q,$$

(2) (Tameness at the starting and ending points)

$$\begin{aligned} \cot \angle(E_{i_1}^u, E_{i_1}^s), \cot \angle(E_{i_2}^u, E_{i_2}^s) &< \frac{D_1^{M\Delta}}{100}, \\ 10^6 \bar{r} &\geq r_i \geq \tau_i, \quad \forall i_1 \leq i \leq i_2 \\ r_{i_1} &= \tau_{i_1} = \bar{r}, \quad \kappa_{i_1} = \tilde{\kappa}_{i_1} = \bar{\kappa}, \\ r_{i_2} &= 10^6 \bar{r}, \quad \tau_{i_2} \leq \frac{1}{10} \bar{r}, \quad \kappa_{i_2} = \frac{1}{100} \bar{\kappa}, \quad \tilde{\kappa}_{i_2} = 100 \bar{\kappa}, \\ \sum_{n=i_1}^{i_2-1} \lambda_n^u, \sum_{n=i_1}^{i_2-1} -\lambda_n^s &\geq \frac{2}{3} (i_2 - i_1) a, \end{aligned}$$

(3) (Transversal mappings) Let r_n, τ_n, κ_n be as above, we let

$$U_n = U(r_n, \tau_n, \kappa_n), \quad C_n = C(\kappa_n), \quad \tilde{C}_n = \tilde{C}(\tilde{\kappa}_n)$$

If Γ is a κ_n -full horizontal graph of U_n , then $g_n(\Gamma) \cap U_{n+1}$ is a κ_{n+1} -full horizontal graph of U_{n+1} . Moreover, the image of the horizontal boundary of U_n under g_n is disjoint from the horizontal boundary of U_{n+1} ; the image of the vertical boundary of U_n under g_n is disjoint from the vertical boundary of U_{n+1} .

(4) (Cone condition) Furthermore, for any $(v, w) \in U_n$, we have $(Dg_n)_{(v,w)}(C_n) \subset C_{n+1}$; for any $(v, w) \in g_n(U_n) \cap U_{n+1}$, we have $(Dg_n^{-1})_{(v,w)}(\tilde{C}_{n+1}) \subset \tilde{C}_n$. Moreover, for any $(v, w) \in U_n$, any $(V, W) \in C_n$, let $(\tilde{V}, \tilde{W}) = (Dg_n)_{(v,w)}(V, W)$, we have $|\tilde{V}| \geq e^{\lambda_n^u - \delta} |V|$; for any $(v, w) \in g_n(U_n) \cap U_{n+1}$, any $(V, W) \in \tilde{C}_{n+1}$, let $(\tilde{V}, \tilde{W}) = (Dg_n^{-1})_{(v,w)}(V, W)$, we have $|\tilde{W}| \geq e^{-\lambda_n^s - \delta} |W|$.

(5) (Hyperbolic map) Denote

$$\begin{aligned} J &= i_{i_1}^{-1} \exp_{x_{i_1}}^{-1} \exp_{x_{i_2}} i_{i_2}, \\ G &= i_{i_1}^{-1} \exp_{x_{i_1}}^{-1} g^{i_2-i_1} \exp_{x_{i_1}} i_{i_1} = J g_{i_2-1} \cdots g_{i_1}. \end{aligned}$$

There exist $\mathcal{R}_1$, a $100\bar{\kappa}$ -vertical strip of U_{i_1} , and $\mathcal{R}_2$, a $100\bar{\kappa}$ -horizontal strip of U_{i_1} such that G is a hyperbolic map from $\mathcal{R}_1$ to $\mathcal{R}_2$ with parameters $\kappa' = \bar{\kappa}, \kappa'' = 100\bar{\kappa}$. Moreover, for each $0 \leq j \leq i_2 - i_1$, we have $g_{i_1+j-1} \cdots g_{i_1}(\mathcal{R}_1) \subset U_{i_1+j}$.

We will give a sketch of the proof and refer the detailed estimates to [2].

Proof. Set $a = \log A_1$. Condition (4) in Proposition 3 translates into

$$\frac{1}{q} \sum_{i=0}^{q-1} \lambda_i^s \leq -a, \quad \frac{1}{q} \sum_{i=0}^{q-1} \lambda_i^u \geq a.$$

Using condition (3) and Lemma 4 in place of the Pliss lemma, by setting $\theta_0 \in (0, 1)$ to be an absolute constant sufficiently close to 1, and setting $C > 0$ to be sufficiently large depending only on X , we can show analogously to Lemma 4.4 in [2], that there are more than $q/2$ points in $\{g^k(x) | 0 \leq k \leq q-1\}$ that are “good in the orbit of x ”. Here a point $g^n(x)$ is said to be *good in the orbit of x* if $n \in [1, q-1]$ satisfies the following conditions :

$$(A.5) \quad \frac{1}{k} \sum_{j=n}^{n+k-1} \bar{\lambda}_j^e > (1 - \frac{1}{1000}) \theta_0^{-1} a, \forall 1 \leq k \leq q-n,$$

$$(A.6) \quad \frac{1}{k} \sum_{j=n-k}^{n-1} \bar{\lambda}_j^e > (1 - \frac{1}{1000}) \theta_0^{-1} a, \forall 1 \leq k \leq n.$$

We can show in analogy to Lemma 4.5 that $|\cot \angle(v_n^s, v_n^u)| \leq A_1^{3\Delta}$ for all n such that $g^n(x)$ is good in the orbit. Then there exist an integer $0 \leq i \leq D_1^{-C_1\Delta} q$ such that the subsequence $(x_{i+j[D_1^{-C_1\Delta} q]})_{0 \leq j \leq \lfloor \frac{q}{[D_1^{-C_1\Delta} q]} \rfloor - 1}$ contains at

least $\frac{1}{3} D_1^{C_1\Delta}$ many points which are good in the orbit of x . By letting C_1 to be sufficiently large depending only on X , we can apply the pigeonhole principle to the above subsequence as in the proof of Proposition 4.1 in [2] and obtain $0 \leq i_1 < i_2 \leq q-1$ that satisfy the following conditions:

- (1) $i_2 - i_1 \geq D_1^{-\Delta C_1} q$,
- (2) $\sum_{j=i_1}^{i_1+k-1} \bar{\lambda}_j^e > (1 - \frac{1}{1000}) \theta_0^{-1} a k, \forall 1 \leq k \leq i_2 - i_1$,
- (3) $\sum_{j=i_2-k}^{i_2-1} \bar{\lambda}_j^e > (1 - \frac{1}{1000}) \theta_0^{-1} a k, \forall 1 \leq k \leq i_2 - i_1$,
- (4) The angles $\angle(v_{i_1}^s, v_{i_1}^u), \angle(v_{i_2}^s, v_{i_2}^u)$ satisfy

$$\log |\cot \angle(v_{i_1}^s, v_{i_2}^u)| \leq 3\Delta a, \quad \log |\cot \angle(v_{i_2}^s, v_{i_2}^u)| \leq 3\Delta a,$$

(5) Moreover, we have $d(g^{i_1}(x), g^{i_2}(x)) < D_1^{-\frac{C_1 \Delta}{200}}$, and

$$d_{T^1 X}(v_{i_1}^s, v_{i_2}^s) < D_1^{-\frac{C_1 \Delta}{200}}, \quad d_{T^1 X}(v_{i_1}^u, v_{i_2}^u) < D_1^{-\frac{C_1 \Delta}{200}}.$$

We note the similarities between the above conditions and those of Definition 4.3 in [2]. However here we have a large inverse power of D_1 in (5) instead of a small inverse power of q as in Definition 4.3, (4) in [2]. This is sufficient for the rest of proof, since r_{i_1}, r_{i_2} , $\angle(E_{i_1}^s, E_{i_1}^u)$ and $\angle(E_{i_2}^s, E_{i_2}^u)$ are lower bounded by $D^{-O(\Delta)}$.

At this point, we can invoke the proof of Proposition 4.2, and obtain (2) as a consequence of Lemma 4.6, 4.7, 4.8 in [2]; and obtain (3),(4) as a consequence of Proposition 4.5 in [2]. We obtain (5) following the proof of Proposition 4.4 in [2]. $\square$

Now we are ready to conclude the proof of Proposition 3.

Proof of Proposition 3. We apply Proposition 6 and obtain $i_1, i_2, \mathcal{R}_1, \mathcal{R}_2, G, U_i, C_i, \tilde{C}_i$ as in the proposition. We set $i = i_1, j = i_2$. By (5) in Proposition 6 and Proposition 4.3 in [2], we obtain a hyperbolic periodic point in $\mathcal{R}_1 \cap \mathcal{R}_2$, denoted by y .

We note the following lemma whose proof follows from the standard construction of unstable / stable manifolds for uniformly hyperbolic maps using graph transform argument. For this reason, we omit its proof.

LEMMA 5. *Let $r, \tau > 0, L > 1, 0 < \kappa, \kappa', \kappa'' < 1, U = U(r, \tau, \kappa)$ and let $G : \mathcal{R}_1 \rightarrow \mathcal{R}_2$ be a hyperbolic map where $\mathcal{R}_1$ (resp. $\mathcal{R}_2$) is the κ -vertical strip (resp. κ -horizontal strip) of U as in Definition 7, and κ', κ'' satisfy inclusion (A.3), (A.4) respectively. Assume that*

(1) *For each $x \in \mathcal{R}_1$, each $(V, W) \in C(\kappa')$, set $(\bar{V}, \bar{W}) = DG_x(V, W)$, then $|\bar{V}| \geq L|V|$,*

(2) *For each $x \in \mathcal{R}_2$, each $(V, W) \in \tilde{C}(\kappa'')$, set $(\bar{V}, \bar{W}) = DG_x^{-1}(V, W)$, then $|\bar{W}| \geq L|W|$.*

Then there exists a hyperbolic fixed point of G , $y \in \mathcal{R}_1 \cap \mathcal{R}_2$, whose local unstable manifold in $\mathcal{R}_2$, denoted by $\mathcal{W}_G^u(y)$, is a κ' -horizontal graph; whose local stable manifold in $\mathcal{R}_1$, denoted by $\mathcal{W}_G^s(y)$, is a κ'' -vertical graph. Moreover we have

$$G(\mathcal{R}_1) \subset B(\mathcal{W}_G^u(y), 2L^{-1} \text{diam}(U)).$$

We set $L = A^{\frac{j-i}{2}}$. We now verify conditions (1),(2) of Lemma 5 for $L, G, U = U_{i_1}, \kappa = 100\bar{\kappa}, \kappa' = \bar{\kappa}, \kappa'' = 100\bar{\kappa}$. We only verify condition (2) in details since condition (1) can be verified in a similar fashion. By Proposition 6(5), for any $i \leq n \leq j-1$, we have $g_{n+1}^{-1} \cdots g_{j-1}^{-1} J^{-1}(\mathcal{R}_2) = g_n \cdots g_i(\mathcal{R}_1) \subset U_{n+1} \cap g_n(U_n)$. For any $i \leq n \leq j$, any $(v, w) \in \mathcal{R}_2$, for any $(V, W) \in \tilde{C}_j$ (here $\tilde{C}_j$ is given by Proposition 6(3)), denote by $(v_n, w_n) = g_n^{-1} \cdots g_{j-1}^{-1} J^{-1}(v, w)$, $(V_n, W_n) = D(Jg_{j-1} \cdots g_n)_{(v,w)}^{-1}(V, W)$. Then we have

$(v_n, w_n) \in U_n$ for all $i \leq n \leq j$. By Proposition 6(2),(4), we have $|W_i| \geq e^{\sum_{n=i}^{j-1} (-\lambda_n^s - \delta)} |W_j| \geq A^{\frac{j-i}{2}} |W| = L|W|$.

By Lemma 5 and Proposition 6(2), we obtain

$$G(\mathcal{R}_1) \subset B(\mathcal{W}_G^u(y), 200A^{-\frac{j-i}{2}} \bar{r})$$

We denote by $z = \exp_{x_{i_1}} i_{i_1}(y)$. By Proposition 6(5) and the fact that y is a hyperbolic fixed point of G , we conclude that z is a g -hyperbolic periodic point. Then by Proposition 6 and by possibly increasing C_1 depending only on X , we can ensure that $z \in \mathcal{H}(g, D_1^{-C_1 \Delta})$, and

$$g^j(x) \in \exp_{x_{i_1}} i_{i_1} G(\mathcal{R}_1) \subset B(\mathcal{W}_{D_1^{-3C_1 \Delta}}^u(z), A_1^{-\frac{q}{2D_1^{C_1 \Delta}}})$$

We conclude the proof by letting C to be sufficiently large depending only on X . □

APPENDIX B.

Proof of Corollary B. In this following, we briefly denote $H(f, \alpha, r)$ by $H(\alpha, r)$, and denote $H(f, \lambda)$ by $H(\lambda)$.

We first prove the corollary under condition (1). For any $\alpha, r, \xi > 0$, any $y \in \mathcal{H}(\alpha, r)$,

$$|B(\mathcal{W}_r^u(y), \xi)| \ll r\xi.$$

It is clear from the definition of $\mathcal{G}$ in (3.1) that for any $\lambda \in (0, 1)$,

$$\begin{aligned} |\mathcal{H}(\lambda)| &\geq |\mathcal{G}(\lambda, \xi, f)| / |B(\mathcal{W}_{\lambda^3}^u(y), \xi)| \\ &\gg \lambda^{-3} \xi^{-1} |\mathcal{G}(\lambda, \xi, f)| \end{aligned}$$

By (4.7) and Proposition 2, it suffices to check that $|\mathcal{G}_i| \gg A^{-\frac{q_{i+1}\theta_0^{i+1}}{2D^{C\Delta q_i}}} D^{11C\Delta q_i}$. Since $x^{-1} > e^{-x}$ for $x \in (0, \infty)$, we have

$$\begin{aligned} |\mathcal{G}_i| &\geq \frac{\eta^{K-i}\varepsilon}{(K+1)l_i} \\ &> (10K)^{-1} \eta^K \varepsilon \frac{q_{i+1}\theta_0^{i+1}}{D^{C\Delta q_i} l_i} A^{-\frac{q_{i+1}\theta_0^{i+1}}{4D^{C\Delta q_i}}} \\ &\gg A^{-\frac{q_{i+1}\theta_0^{i+1}}{2D^{C\Delta q_i}}} D^{11C\Delta q_i}. \end{aligned}$$

The last inequality follows from by letting $C' \gg 1$, and

- $K^{-1} \eta^K \varepsilon q_{i+1} \theta_0^{i+1} l_i^{-1} \gg 1$ since by (4.4), and $q_{i+1} l_i^{-1} \theta_0^{i+1} \geq \frac{1}{2} q_i \theta_0^i \geq \frac{1}{2} q_0 \geq \frac{1}{2} \varepsilon^{-1} e^{C'(\log(\frac{\log A}{h}))^2 + C'} \gg \varepsilon^{-1} K \eta^{-K}$,
- $A^{\frac{q_{i+1}\theta_0^{i+1}}{4D^{C\Delta q_i}}} \geq D^{12C\Delta q_i}$, by $A \gg 1$, (4.3) and (4.5).

Now we consider condition (2). We set $\delta_0 = D^{-Tower(Q_0, Q_1, K+3)}$. By Lemma 1(3), we have

$$\delta_0 \leq \xi_0 \text{ and } \delta_0 \leq \lambda_i^3, \forall 0 \leq i \leq K.$$

For any $\lambda, \xi \in (0, 1)$, any $y \in \mathcal{H}(\lambda)$, any $\delta \in (0, \lambda^3)$, we have

$$m(B(B(\mathcal{W}_{\lambda^3}^u(y), \xi), \delta)) \ll \lambda^3 \max(\xi, \delta).$$

By (3.1) and condition (2), we have for some $0 \leq i \leq K$ that,

$$\begin{aligned} |\mathcal{H}(\lambda_i)| &\geq m(B(\mathcal{G}(\lambda_i, \xi_i, f), \delta_0)) / \sup_{y \in H(\lambda_i)} m(B(B(\mathcal{W}_{\lambda_i^3}^u(y), \xi_i), \delta_0)) \\ &\gg \varepsilon \lambda_i^{-3} \min(\xi_i^{-1}, \delta_0^{-1}). \end{aligned}$$

By Proposition 2, it suffices to observe from Lemma 1 that

$$\varepsilon \gg \lambda_i^{-11} \max(\xi_i, \delta_0), \quad \forall 0 \leq i \leq K.$$

□

REFERENCES

- [1] L. M. Abramov, V. A. Rohlin, *The entropy of a skew product of measure preserving transformations*, AMS Translations, 48 (1965), 255-265.
- [2] A. Avila, B. Fayad, P. Le Calvez, D. Xu, Z. Zhang, *On mixing diffeomorphisms of the disk*, arXiv:1509.06906.
- [3] V. Climenhaga, D. Dolgopyat, Y. Pesin, *Non-stationary non-uniform hyperbolicity: SRB measures for non-uniformly hyperbolic attractors*, Communications in Mathematical Physics, 346, issue 2 (2016), 553-602.
- [4] V. Climenhaga, Y. Pesin, *Hadamard-Perron theorems and effective hyperbolicity*, Ergodic Theory and Dynamical Systems, 36 (2016), 23-63.
- [5] A. Katok, *Lyapunov exponents, entropy and periodic orbits for diffeomorphisms*, Publications Mathématiques de l'IHÉS, 51 (1980), 137-174.
- [6] A. Katok, B. Hasselblatt, *Introduction to the Modern Theory of Dynamical Systems*. Encyclopedia of Mathematics and its Applications 54. Cambridge: Cambridge University Press, 1995.
- [7] V. Pliss, *On a conjecture of Smale*, Diff. Uravnenjia, 8:268-282, 1972.

Bassam Fayad
 Institut de Mathématiques de Jussieu-Paris Rive Gauche UMR7586 CNRS Université Paris Diderot-Université Pierre et Marie Curie
 E-mail: bassam.fayad@imj-prg.fr

Zhiyuan Zhang
 Institut de Mathématique de Jussieu—Paris Rive Gauche, Bâtiment Sophie Germain, Bureau 652
 75205 PARIS CEDEX 13, FRANCE
 Email address: zzzhangzhiyuan@gmail.com